

## **Adult New Patient Intake Form**

Patient Informa	tion					
Last Name:			me:			DOB:
Gender: H	lome Phone:			Mobile F	Pho	one:
Preferred Phone: Home or Mobile (circle one)			Email: _			
Emergency Contact:				Relation	ish	ıp:
Emergency Con	act Phone:			Patient	Ma	arital Status:
Occupation:				Employe	er:	
Primary Care Pr					PCP Phone:	
Referring Provid	er:				_	
Preferred Pharm	acy:				_	Pharm Phone:
Preferred Pharm	acy Address:					
Please list ALL a	ctive treating	physicians (i.e. pulm	nonologi	st, oncolo	gis	st, internist, cardiologist, etc)
Doctor's Name:			Speci	alty:		
Doctor's Name:			Speci	alty:		
Doctor's Name:			Speci	alty:		
	_	ormation is encourag	_		altl	h agencies. This information is used to
Ethnicity:	R	ace:				
□ Decline Respo	nse 🗆	Decline Response			_	Black or African American
		American-Indian or A	Alaska N	ative	_	Native Hawaiian or Pacific Islander
□ Not Hispanic o	r Latino 🗆	Asian				White   Other
Preferred Langua	ge:					Decline Response
responsible and r benefits be paid or release pertinent	all applicable on the full payment directly to Columedical inform	opayments and deducent for all charges not on the mbia Doctors for servicention to my insurance	covered l ces rende compan	by my insu red. I auth y when re	ırar nori	ime of service. I agree to be financially nce company. I authorize my insurance ize representatives of ColumbiaDoctors to ested or to facilitate payment of a claim.
I acknowledge th  ☐ Received ☐ N	at I was provido /A (only if you	received the notice fro	Columbial	Doctors No		ce of Privacy Practices (NOPP). reviously)
	will provide yours	ou with the health plan	•	•		a) accepts*. If you decide to be treated by a consent form agreeing that you accept
I read and agree	to all of the ab	ove (Financial Agreem	nent, No	tice of Priv	vac	ry, Insurance Information).
Patient or Leg	al Guardian Na	me (Print):				
Patient or Leg	al Guardian Sig	nature:				Date:

\*Please refer to our website: columbiadoctors.org, for a list of insurances accepted by your provider.

# COLUMBIA UNIVERSITY HEADACHE CENTER: NEW PATIENT QUESTIONNAIRE

## **HEADACHE CHARACTERISTICS**

Frequency and Seve	erity							
1. At what AGE did you get your <u>first</u> headache, of ANY kind?								
2. At what AGE did your headaches become problematic?								
3. On <u>AVERAGE</u> , how many DA	AYS per MONTH do y	ou get hea	adaches, of	ANY KIND?				
4. On <u>AVERAGE</u> , how long do	your typical headach	es last? (N	AINUTES, H	OURS, DAYS)				
5. On a scale of zero to ten, ho	ow would you rate yo	our <u>WORS</u>	T pain?		(please	e circle your answe	r)	
0 1	2 3 4	5	6	7	8	9 10		
no pain	mild -	mode			severe	5 10		
6. On a scale of zero to ten, ho	ow would you rate yo	ur <u>AVERA</u>	GE pain?		(please	e circle your answe	r)	
0 1	2 3 4	5	6	7	8	9 10		
no pain	z 3 4 mild	mode			severe	9 10		
7. How many DAYS per MONT	H do you get your <u>W</u>	ORST hea	daches?					
8. How long do your WORST h	eadaches last? (MIN	UTES, HO	URS, DAYS)					
Location(s)								
	Left	R	Right		Front		Back	
Please indicate the regions affected:								
2. Is one side affected more than another?	Left side only Yes,% more on the left side	Υ	Right side or es, nore on the	_%		th sides ally affected	No, alternates equally	
3. What region(s) is/are affected?	☐ Front of the head ☐ Back of the head ☐ Side of the head ☐ Top of the head	I	Around the Ey Behind the Eye Shoulder Neck			☐ Ear ☐ Nose ☐ Jaw ☐ Other		
Quality/Description of Pain								
How would you describe your	_			The state of the s	e (∜)chec	k ALL that apply:		
☐ Throbbing ☐ Tearing	☐ Stabbing ☐ Sharp		_	ounding		☐ Pulsating ☐ Burning		
Dull	☐ Sharp ☐ Pressure ☐ Gradual					Constant		
Aching	☐ Squeezing		□ т	ightening		Other		
Associated Feature	s							
When you have head pain, do	you have any of the	following	?	Pleas	se (√)che	ck ALL that apply:		
Sensitivity to light		Rooms	-		<del></del>	Tearing of the eye	( right / left / both )	
Sensitivity to sound		Self spi	_		_	Redness of the eye		
Sensitivity to smell		☐ Lighthe	eadedness		_	Drooping of the ey Nasal congestion	e (right / left / both)	
Sensitivity to movement			vision ty concentr	ating		Runny nose	( right / left / both ) ( right / left / both )	
☐ Nausea ☐ Vomiting			,	. 0	_	Facial flushing	( right / left / both )	
⊔ *∨						For fullmoss	/ right / loft / both \	

☐ Ear fullness

( right / left / both )

Have you experienced any	of the fo	llowing?		Please (√)check ALL that	apply:				
Head or neck trauma?		□ No		·					
			describe:						
Loss of consciousness?		☐ No ☐ Yes, describe:							
Infection/illness BEFORE th	ne	□No							
onset of your headache	e?		describe:						
Toxic exposure BEFORE the	e onset	☐ No							
of your headache?			describe:						
Snoring or breathing pause	es?	☐ No ☐ Yes, o	describe:						
Jaw clicking, difficulty oper	ning	☐ No							
mouth, tooth grinding?		☐ Yes, o	describe:						
Use of tetracycline antibio									
retinoic acid/vitamin A		□ No							
BEFORE the onset of you headache?	our	∐ Yes, o	describe:						
Headache:									
Have you been diagnosed	with any	of the fo	llowing?	Please (√)check ALL that	apply:				
Anxiety		polar Dis		Depression	☐ Other psychiatric disorder				
Anorexia		ılimia	order	☐ Posttraumatic Stress	☐ Other psychiatric disorder				
Psychosis	_		Compulsive Disorder	_ r ostadamade sa ess					
_ ,	_		•						
DIAGNOSTIC ST					ale and a fit and a second a fit and a fit a				
Study	Da		Findings	at have been performed: (Atta	ch copy of test results if possible)				
•	Da	ite	Findings						
BRAIN MRI									
NECK (Cervical) MRI									
MRA/MRV									
Head CT									
Lumbar puncture									
Sleep Study									
EEG									
Other									

### **PAST PREVENTIVE MEDICATIONS**

PASI PREVI	ENTIVE MEDICA			8:14: 1 4	
Medication Generic Name ( <i>Brand Name</i> )	What was the highest dose (mg)?	How many times per day?	When/how long was this used?	Did this change the intensity, duration or frequency of the headache? (If so, how?)	Were there any side effects? (If so, what were they?)
Amitriptyline ( <i>Elavil</i> )					
Nortriptyline ( <i>Pamelor</i> )					
Doxepin (Sinequan)					
Imipramine (Tofranil)					
Cyproheptadine (Periactin)					
Venlafaxine (Effexor)					
Duloxetine (Cymbalta)					
Escitalopram (Lexapro)					
Topiramate ( <i>Topamax</i> )					
Zonisamide (Zonegran)					
Valproate ( <i>Depakote</i> )					
Gabapentin (Neurontin)					
Lamotrigine (Lamictal)					
Carbamazepine (Tegretol)					
Oxcarbazepine (Trileptal)					
Acetazolamide ( <i>Diamox</i> )					
Propranolol (Inderal)					
Nadolol (Corgard)					
Atenolol (Tenormin)					
Metoprolol (Toprol)					
Candesartan (Atacand)					
Flunarizine (Sibelium)					
Verapamil (Calan, Veralan)					
Indomethacin (Indocin)					
Lithium (Eskalith, Lithobid))					
Methysergide (Sansert)					
Vitamin B2 ( <i>Riboflavin</i> )					
Magnesium					
Butterbur (Petadolex)					
Feverfew					
Melatonin					

#### PAST ACUTE MEDICATIONS

PAST ACUTE ME	Dose (mg)	Route of administration (oral, nasal, injection)	Did this change the intensity or duration of the headache? (If so, how?)	Were there any side effects? (If so, what were they?)
Sumatriptan ( <i>Imitrex</i> )				
Treximet				
Rizatriptan (Maxalt)				
Zolmitriptan ( <i>Zomig</i> )				
Naratriptan ( <i>Amerge</i> )				
Eletriptan ( <i>Relpax</i> )				
Almotriptan (Axert)				
Frovatriptan ( <i>Frova</i> )				
Ergotamine tablets ( <i>Cafergot, Ergomar, Migergot, Migracet, Wigraine</i> )				
Ergotamine suppository (Cafergot)				
Dihydroergotamine nasal spray (Migranal)				
Promethazine ( <i>Phenergan</i> )				
Chlorpromazine ( <i>Thorazine</i> )				
Prochlorperazine (Compazine)				
Metoclopramide (Reglan)				
Naproxen (Naprosyn, Aleve)				
Ibuprofen (Advil, Motrin)				
Acetaminophen/Aspirin/Caffeine (Excedrin)				
Nabumetone ( <i>Relafen</i> )				
Flurbiprofen ( <i>Ansaid</i> )				
Mefenamic Acid (Ponstel)				
Ketorolac ( <i>Toradol</i> )				
Diclofenac (Cambia, Voltaren)				
Steroids (e.g., Medrol, prednisone)				
Muscle relaxants (e.g., tizanidine, methocarbamol)				
Butalbital-containing compounds (Fioricet, Fiorinal)				

## IN THE PAST, WHAT OTHER MEDICATIONS FOR HEADACHE HAVE YOU USED? (Attach additional sheets if needed)

Medication	What was the highest dose (mg)?	How many times per day?	When/how long was this used?	Did this change the intensity, duration or frequency of the headache? (If so, how?)	Were there any side effects? (If so, what were they?)

## INTRAVENOUS THERAPIES, HOSPITALIZATIONS, and PROCEDURES FOR HEADACHE

Treatment	Date	Description of Treatment and Effect
Intravenous DHE (dihydroergotamine)		
Intravenous valproic acid (Depakote)		
Intravenous lidocaine		
Greater occipital nerve block		
Botox Injections		
Implanted stimulator device		
Surgical intervention		
Other:		

### **ADJUNCTIVE THERAPIES**

Type of Treatment	Date	Effect
Cefaly (supraorbital nerve stimulator)		
Transcranial Magnetic Stimulation (TMS)		
Oxygen therapy		
Acupuncture		
Biofeedback		
TENS therapy		
Chiropractic Manipulation		
Massage Therapy		
Physiotherapy		
Homeopathy		
Other:		

#### **MIDAS (Migraine Disability Assessment)**

#### About MIDAS:

The MIDAS (migraine disability assessment) questionnaire is a tool to help you measure the impact your headaches have on your life over the last 3 months. The best way to this is by counting the numbers of days of your life which are affected by Headaches. You can do this for yourself as follows:

#### **INSTRUCTIONS:**

- Please complete Questions about ALL your headaches you have had over the last 3 months.
- Write your answer in the box next to each question.
- If a single headache affects more than one area of your life (e.g., work and family life) it is counted more than once.
- Mark question as zero if you did not do the activity in the last three months.
- · For questions 1 and 2, work or school means paid work or education if you are a student at school or college.
- For Questions 3 and 4, **household work** means activities such as housework, home repairs and maintenance, shopping as well as caring for children and relatives.

1. On how many days in the last 3 months did you miss work or school because of headaches?	days	
2. How many days in the last 3 months was your productivity at work or school r more because of your headaches? (Do not include days you counted in quest missed work or school)	days	
3. On how many days in the last 3 months did you not do household work becau headaches?	days	
4. How many days in the last 3 months was your productivity in household work more because of your headaches? (Do not include days you counted in quest did not do household work)	days	
5. On how many days in the last 3 months did you miss family, social or leisure a of your headaches?	days	
To score, add points for answers in each column	Total:	days

MIDAS	MIDAS	Definition	Recommendations
Score 0–5	Grade I	Minimal or infrequent disability	MIDAS Grade I usually indicates low medical need. Simple, over-the-counter analgesics may be effective in the acute treatment of these patients. However, the impact of even a few lost days on the lifestyle of these patients should be assessed. Also, some patients with a MIDAS Grade I, such as those with infrequent, but severe migraine, may benefit from first-line treatment with specific migraine therapies (e.g. triptans). MIDAS Grade I patients who have failed to achieve effective relief with simple analgesics should also be considered for triptan therapy.
6–10	II	Mild or infrequent disability	MIDAS Grade II usually indicates moderate medical need. The patients may require an acute prescription medication. Some MIDAS Grade II patients may also qualify for first-line triptan medication if their headaches are severe. For example, a score of 10 could mean that a patient is missing ten days of school or paid work, so the headaches could be causing severe disruption in their lives. MIDAS Grade II patients should also qualify for first-line triptan medication if they have failed on simple analgesics.
11–20	III	Moderate disability	MIDAS Grade III/IV indicates a high medical need. These patients are experiencing significant disability and their migraine attacks are having a severe impact on their lives. Specific acute therapy, such as a triptan, is usually the most appropriate therapy for these patients,
≥21	IV	Severe disability	providing they are suitable recipients. Prophylactic treatment should also be considered.  Please note that a very high MIDAS score could also indicate a high frequency of non- migraine headache, and these patients should be managed accordingly.

Patient's Signature	
Reviewed with the patient on	
Physician's Signature	